

# PHOTOSYNTHETIC PERFORMANCE AND PRODUCTIVITY OF PHYTOPLANKTON IN THE SOUTHERN OCEAN

by

Wee Cheah, B.Sc. Hons, M.Sc.

Submitted in fulfilment of the requirements  
for the degree of Doctor of Philosophy

Institute for Marine and Antarctic Studies  
University of Tasmania  
September, 2012



I declare that this thesis contains no material which has been accepted for a degree or diploma by the University or any other institution, except by way of background information and duly acknowledged in the thesis, and that, to the best of my knowledge and belief, this thesis contains no material previously published or written by another person, except where due acknowledgement is made in the text of the thesis, nor does the thesis contains any material that infringes copyright.

Signed: \_\_\_\_\_  
Wee Cheah

Date: \_\_\_\_\_

This thesis may be reproduced, archived, and communicated in any material form in whole or in part by the University of Tasmania or its agents, and made be made available for loan and copying in accordance with the *Copyright Act 1968*.

Signed: \_\_\_\_\_  
Wee Cheah

Date: \_\_\_\_\_

## Abstract

Marine phytoplankton account for approximately half of global primary production, an amount equivalent to their terrestrial counterpart. These short-lived organisms, with turnover rates between one and three weeks, support nearly all life in the ocean and have a profound effect on global biogeochemical cycles and climate. The connection between marine phytoplankton and climate is intimate and changes to either will profoundly affect the other. Over the years, due to high operational costs and distance from major human settlements, the Southern Ocean has been the least studied ocean, despite its significance in the distribution of nutrients to the world oceans, especially the lower latitudes, and controlling global climate. In order to capture the response of the phytoplankton to environmental change across the vast Southern Ocean, a method with high spatio-temporal resolution is desirable. By focusing on the Australian sector of the Southern Ocean, this dissertation examines the productivity and physiology of natural phytoplankton communities *in situ* using the fast repetition rate (FRR) fluorometry technique.

The FRR fluorometry technique was used to derive direct estimation of *in situ* primary productivity in the Southern Ocean during the SAZ-Sense (Sub-Antarctic Zone Sensitivity to Environmental Change) voyage in Jan-Feb 2007. A statistically significant correlation between FRR- and  $^{14}\text{C}$ -derived primary production was observed ( $r^2 = 0.85$ , slope =  $1.23 \pm 0.05$ ,  $p < 0.01$ ,  $n = 85$ ) but the relationship between the methods differed vertically and spatially, mainly due to the effect of non-photochemical quenching under high irradiance. This indicates the FRR fluorometry technique can be used to determine *in situ* primary productivity in the Southern Ocean but care should be taken in the interpretation of the data.

In addition to the primary production measurements, the photosynthetic performance of phytoplankton was investigated to provide a better understanding of how natural phytoplankton communities acclimate to different environmental variables, especially in the iron-replete Subantarctic Zone (SAZ) and iron-depleted Polar Frontal Zone (PFZ). High effective

photochemical efficiency of photosystem II ( $F'_q/F'_m > 0.4$ ), maximum photosynthesis rate ( $P_{\max}^B$ ), light-saturation intensity ( $E_k$ ), maximum rate of photosynthetic electron transport ( $1/\tau_{\text{PSII}}$ ), and low photoprotective pigment concentrations observed in the SAZ correspond to high chlorophyll *a* and iron concentrations. In contrast, phytoplankton in the PFZ exhibits low  $F'_q/F'_m$  ( $\sim 0.2$ ) and high concentrations of photoprotective pigments under low light environment. Strong negative relationships between iron, temperature, and photoprotective pigments demonstrate that cells were producing more photoprotective pigments under low temperature and iron conditions, and are responsible for the low biomass and low productivity measured in the PFZ.

FRR fluorometry data from 31 transects collected aboard MV I'Astrolabe between 2002 and 2009, were used to assess the photosynthetic performance of phytoplankton along a repeated transect from Hobart (42.8°S, 147.3°E) to the French Antarctic station, Dumont d'Urville (66°S, 140°E). The maximum photochemical efficiency of photosystem II ( $F_v/F_m$ ) values were high in the Subtropical Zone and water close to the Antarctic continent, but low in the PFZ. Spring  $F_v/F_m$  were higher than other seasons, suggesting higher nutrient supply. High  $F_v/F_m$  observed in the Subtropical Zone and Antarctic Zone is consistent with moderate to high iron concentrations in these regions. Overall, phytoplankton photophysiology in the Southern Ocean is governed by nutrient distributions, especially iron, which are affected by atmospheric and oceanic physical processes.

## Acknowledgements

This thesis would not be realized without the supports and helps from many individuals. First, I would like to thank my supervisors, Andrew McMinn and Klaus Meiners for all the scientific conversations over the years, their untiring support throughout this project, and assistance with the thesis preparation. I am truly indebted to my primary supervisor, Andrew McMinn, for his expert advice, sense of humour, dedication, and financial support. I really enjoy and appreciate his supervision and guidance.

Thanks to the officers and crews of RSV Aurora Australis, MV l’Astrolabe, Australian Antarctic Divisions (AAD) gear officers, and fellow expeditioners and volunteers for their support during my three amazing Antarctic experiences. I acknowledge Mark Rosenberg for the oceanography data, Neil Johnson, Alicia Navidad, John Akl, Kristina Paterson for the macronutrients data, Andrew Bowie and Delphine Lannuzel for making the iron data available, Sergei Sokolov and Steve Rintoul for the oceanographic fronts data.

Thanks to all staff and students at the Institute of Antarctic and Southern Ocean Studies (IASOS), Institute for Marine and Antarctic Studies (IMAS), and Antarctic Climate and Ecosystems Cooperative Research Centre (ACE CRC) for providing such a great working atmosphere. In particular, thanks to Margaret Hazelwood, Juliar Jabour, Patti Virtue, Kelvin Michael, Kate Maloney, Denbeigh Armstrong, Jemina Stuart-Smith, Bernadette Ulbrich-Hooper for their their assistance, and Rob Johnson, Pier van der Merwe, Alex Fraser, Miguel de Salas, Helena Baird, Tom Remenyi, Alyce Hancock, Jake van Oosterom, Ming Fu, Sazlina Salleh, Mahadi Mohammad, Leonie Jordan, Krystyna Saunders, Andrew Martin, Fraser Kenney, Laura Smith, Simon Geard, and Simon Reeves, for the many conversations on phytoplankton, oceanography, fluorometers, and computing.

I would like to thank Brian Griffiths (CSIRO), Simon Wright (AAD), Peter Strutton (IMAS), Karen Westwood (AAD), and Lesley Clementson (CSIRO), for their discussion and contribution to my manuscripts.

I would also like to thank Tony Fenton and his parents (Geoff and Janet), Glenda Ashmore, my landlord Michael Fenton, for their company and kindness, making my life in Tassie such a memorable one. To my Ariba Amoebas underwater team mates (Rob, Yumi, Alex, Dave, Duncan, Gale, Simon, Simon, Sarah, Alyce, Chris, Jake, Amelie, Dachlen, Michael), thanks for the wonderful games.

Finally, I would like to dedicate this thesis to my family and Hiroko for their understanding and encouragement over the years.

This work was supported financially by the Australian Government through the Australian Antarctic Science Grants Project (#2720 and #1307) and ACE CRC. I would also like to thank ACE CRC for providing me a PhD scholarship.

# **The following Chapters have been removed for copyright or proprietary reasons**

## **Chapter 1.**

W. Cheah, A. McMinn, F. B. Griffiths, K. J. Westwood, J. P. Webb, E. Molina, S. W. Wright, and R. van den Enden. 2011. Assessing Sub-Antarctic Zone primary productivity from fast repetition rate fluorometry, *Deep Sea Research II*, 58: 2179-2188. doi: 10.1016/j.dsr2.2011.05.023.

## **Chapter 2.**

W. Cheah, A. McMinn, F. B. Griffiths, K. J. Westwood, S. W. Wright, and L. A. Clementson. Dynamic influences of iron and temperature on phytoplankton photophysiology in the changing Sub-Antarctic Zone, *PLOS ONE*, in revision.

## **Chapter 3.**

W. Cheah, A. McMinn, P. G. Strutton, and F. B. Griffiths. Seasonal to interannual variability in phytoplankton photosynthetic performance in the Southern Ocean, 2002-2009, *Journal of Plankton Research*, in revision.



## Publications resulting from research completed during candidature

### Published/in revision

### Refereed:

1. **W. Cheah**, A. McMinn, F. B. Griffiths, K. J. Westwood, J. P. Webb, E. Molina, S. W. Wright, and R. van den Enden. 2011. Assessing Sub-Antarctic Zone primary productivity from fast repetition rate fluorometry, *Deep Sea Research II*, 58: 2179-2188. *doi*: 10.1016/j.dsr2.2011.05.023.
2. **W. Cheah**, A. McMinn, F. B. Griffiths, K. J. Westwood, S. W. Wright, and L. A. Clementson. Dynamic influences of iron and temperature on phytoplankton photo-physiology in the changing Sub-Antarctic Zone, *PLOS ONE*, *in revision*.
3. **W. Cheah**, A. McMinn, P. G. Strutton, and F. B. Griffiths. Seasonal to interannual variability in phytoplankton photosynthetic performance in the Southern Ocean, 2002-2009, *Journal of Plankton Research*, *in revision*.

## Statement of Co-Authorship

The following people contributed to the publication of the work undertaken as part of this thesis:

Paper 1/Chapter 2 (*Assessing Sub-Antarctic Zone primary productivity from fast repetition rate fluorometry*):

- **Wee Cheah** (81%)
- Andrew McMinn (6%)
- F. Brian Griffiths (4%)
- Karen J. Westwood (3%)
- Simon W. Wright (3%)
- Ernesto Molina (1%)
- Jason P. Webb (1%)
- Rick van den Enden (1%)

Paper 2/Chapter 3 (*Dynamic influences of iron and temperature on phytoplankton photophysiology in the changing Sub-Antarctic Zone*):

- **Wee Cheah** (81%)
- Andrew McMinn (6%)
- F. Brian Griffiths (4%)
- Karen J. Westwood (3%)
- Simon W. Wright (3%)
- Lesley A. Clementson (3%)

Paper 3/Chapter 4 (*Seasonal to interannual variability in phytoplankton photosynthetic performance in the Southern Ocean, 2002-2009*):

- **Wee Cheah** (82%)
- Andrew McMinn (8%)
- Peter G. Strutton (6%)
- F. Brian Griffiths (4%)

## Details of the authors' roles:

Wee Cheah designed and implemented the experiment, performed data analysis and manuscript writing.

Andrew McMinn contributed with project development and refinement, technical and conceptual discussion, and document preparation.

Brian Griffiths and Peter Strutton assisted with technical and conceptual aspects of the papers, as well as document preparation.

Karen Westwood, Simon Wright, and Lesley Clementson provided data and assisted in document preparation.

Ernesto Molina, Jason Webb, and Rick van den Enden provided assistance in data collection.

We the undersigned agree with the above stated proportion of work undertaken for each of the above published (or submitted) peer-reviewed manuscripts contributing to this thesis:

Signed: \_\_\_\_\_

Andrew McMinn

Supervisor

Institute for Marine and Antarctic Studies

Date: \_\_\_\_\_

Signed: \_\_\_\_\_

Mike Coffin

Head of School

Institute for Marine and Antarctic Studies

Date: \_\_\_\_\_

## Table of Contents

<b>Table of Contents</b>	<b>xi</b>
<b>List of Tables</b>	<b>xiii</b>
<b>List of Figures</b>	<b>xiv</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Marine primary producers . . . . .	1
1.2 The Southern Ocean . . . . .	2
1.3 Measuring primary production . . . . .	3
1.4 Fast repetition rate fluorometry . . . . .	4
1.5 Thesis objectives . . . . .	7
<b>2 Assessing Subantarctic Zone primary productivity from fast repetition rate fluorometry</b>	<b>8</b>
2.1 Abstract . . . . .	8
2.2 Introduction . . . . .	9
2.3 Materials and methods . . . . .	11
2.3.1 Sampling . . . . .	11
2.3.2 FRR measurements . . . . .	14
2.3.3 $^{14}\text{C}$ measurements . . . . .	14
2.3.4 FRR primary productivity model . . . . .	16
2.3.5 Data analysis . . . . .	17
2.4 Results . . . . .	17
2.4.1 Hydrological conditions . . . . .	17
2.4.2 Variable fluorescence measurements . . . . .	18
2.4.3 Chlorophyll <i>a</i> , productivity and macronutrients . . . . .	18
2.4.4 FRR and $^{14}\text{C}$ primary productivity estimates . . . . .	21
2.5 Discussion . . . . .	21
2.5.1 Variability of fluorescence measurements . . . . .	21
2.5.2 Primary productivity SAZ-West, SAZ-East and PFZ . . . . .	23
2.5.3 FRR- and $^{14}\text{C}$ -based productivity . . . . .	25
2.6 Conclusion . . . . .	28
<b>3 Dynamic influences of iron and temperature on phytoplankton photophysiology in the changing Sub-Antarctic Zone</b>	<b>30</b>

3.1	Abstract . . . . .	30
3.2	Introduction . . . . .	31
3.3	Materials and Methods . . . . .	33
3.3.1	Hydrography . . . . .	33
3.3.2	FRR Fluorometry . . . . .	34
3.3.3	Photosynthesis-Irradiance Incubations . . . . .	35
3.3.4	Spectral Absorption and Pigments . . . . .	36
3.3.5	Statistics . . . . .	37
3.4	Results . . . . .	37
3.4.1	Hydrographic Conditions . . . . .	37
3.4.2	Photosynthetic Parameters . . . . .	38
3.4.3	Phytoplankton Pigment, Absorption and Community Structure . . . . .	42
3.5	Discussion . . . . .	46
3.5.1	Photoacclimation of Phytoplankton in the SAZ and PFZ . . . . .	46
3.5.2	Influence of Iron and Temperature on Photoacclimation . . . . .	48
<b>4</b>	<b>Seasonal to interannual variability in phytoplankton photosynthetic performance in the Southern Ocean, 2002-2009</b>	<b>53</b>
4.1	Abstract . . . . .	53
4.2	Introduction . . . . .	53
4.3	Data and analysis . . . . .	55
4.4	Results . . . . .	58
4.4.1	Hydrographic properties . . . . .	58
4.4.2	Trends in $F_v/F_m$ . . . . .	59
4.4.3	Diurnal $F_v/F_m$ patterns . . . . .	62
4.4.4	The link between $F_v/F_m$ and environmental variables . . . . .	64
4.5	Discussion . . . . .	66
4.5.1	Trends in $F_v/F_m$ . . . . .	66
4.5.2	Diurnal $F_v/F_m$ patterns . . . . .	69
4.5.3	$F_v/F_m$ and environmental variables . . . . .	70
4.6	Conclusions . . . . .	71
<b>5</b>	<b>Overall summary and conclusion</b>	<b>72</b>
5.1	Summary . . . . .	72
	<b>Bibliography</b>	<b>74</b>

## List of Tables

2.1	Station locations and characteristics. SAZ, Subantarctic Zone; STF, Subtropical Front; PFZ, Polar Frontal Zone; T1, Transit 1; P1, Process 1; P2, Process 2; T3, Transit 3; P3, Process 3; T4, Transit 4 . . . . .	13
2.2	Parameters and definitions used in this study. . . . .	15
3.1	Photosynthetic parameters and definitions used in this study. . . . .	35
3.2	Kendall's rank correlations between bio-optical properties of phytoplankton and environmental variables observed at all stations in this study. Significant correlations at 95 % significance level are indicate in <b>bold</b> . . . . .	41
3.3	Chlorophyll <i>a</i> concentrations ( $\mu\text{g L}^{-1}$ ) and accessory pigment composition normalized to chl <i>a</i> (denoted with asterisks) with depth at each station. Shown are mean (standard deviation). . . . .	44
3.4	Contribution (%) of each phytoplankton group to the total chlorophyll integrated to 120 m at each station. . . . .	46
4.1	Correlations between $F_v/F_m$ and environmental variables in different zones. SAM, Southern Annular Mode; MEI, Multivariate El Niño/Southern Oscillation Index. Significant correlation coefficients are in <b>bold</b> with ** indicates significant level when $p < 0.01$ and * indicates significant level when $p < 0.05$ . . . . .	66
4.2	Correlations between SAM Index, MEI and environmental variables in different zones. SAM, Southern Annular Mode; MEI, Multivariate El Niño/Southern Oscillation Index. Significant correlation coefficients are in <b>bold</b> with ** indicates significant level when $p < 0.01$ and * indicates significant level when $p < 0.05$ . . . . .	67

## List of Figures

1.1	Fast repetition rate fluorometer during deployment mode and measuring technique. The dual (light and dark) optical chambers contain series of blue light-emitting-diodes that provide subsaturating excitation flashes at 200 kHz repetition rate and fluorescence released from excited chlorophyll pigments will be detected by highly sensitive photomultiplier tube. . . . .	4
1.2	Simplified illustration of photosystem II. Light-harvesting pigments such as chlorophyll get excited after absorbing light energy (flashes). The absorbed light energy can then migrate to a reaction centre. If the reaction centre is open, minimum fluorescence $F_o$ is released. If, at the instant of the photon absorption, the reaction centre is closed, the absorbed energy can be reradiated as fluorescence, increasing the fluorescence yield to $F_m$ . Modified from ( <i>Kolber and Falkowski, 1993</i> ) . . . . .	5
1.3	Excitation flashes and relative fluorescence emission during a fast repetition rate fluorometry protocol. The excitation flashes consist of 100 subsaturation flashlets at 1.1 $\mu$ s flash duration and 2.8 $\mu$ s inter flash period. . . . .	6
2.1	Map of the SAZ-Sense sampling area and CTD stations sampled for FRR and $^{14}\text{C}$ measurements. Approximate fronts locations are shown. P1, Process station 1; P2, Process station 2; P3, Process station 3; STF, Subtropical Front; SAF-N, Sub-Antarctic Front North; SAF-S, Sub-Antarctic Front; PF, Polar Front. . . . .	11
2.2	Vertical profiles of $F'_v/F'_m$ , $F'_q/F'_m$ , $q_P$ , $\sigma'_{\text{PSII}}$ and $E$ obtained from two FRR casts at Station 4. (a) and (b) were data from CTD 12 collected at dawn while (c) and (d) were midday measurements from CTD 18. . . . .	19
2.3	Vertical profiles of chl $a$ , FRR-based PP, photochemical efficiency and macronutrients at the three process stations. P1, Process station 1; P2, Process station 2; P3, Process station 3. . . . .	20
2.4	Comparison between primary productivity measured by FRR and $^{14}\text{C}$ for (a) all stations ( $r^2 = 0.85$ , slope = $1.23 \pm 0.05$ s.d., $n = 85$ , $p < 0.01$ ), (b) under light limitation conditions ( $r^2 = 0.96$ , slope = $1.05 \pm 0.03$ s.d., $n = 71$ , $p < 0.01$ ) and (c) under light saturation conditions ( $r^2 = 0.64$ , slope = $1.43 \pm 0.23$ s.d., $n = 14$ , $p < 0.01$ ). The dashed line represents the 1:1 relationship. T1, Transit 1; P1, Process 1; P2, Process 2; T3, Transit 3; P3, Process 3; T4, Transit 4. . . . .	22
2.5	Correlation between the (a) chl $a$ -specific rate of electron transfer ( $ETR^{\text{chl}}$ ) and the carbon uptake rate per unit chl $a$ ( $P_B$ ) for all stations ( $r^2 = 0.74$ , slope = $5.07 \pm 0.28$ s.d., $n = 85$ , $p < 0.01$ ), (b) FRR- and $^{14}\text{C}$ -derived primary productivity normalized by light for all stations ( $r^2 = 0.86$ , slope = $1.03 \pm 0.00$ s.d., $n = 85$ , $p < 0.01$ ). The dashed line represents the 1:1 relationship. . . .	23

2.6	Relationship between FRR- and $^{14}\text{C}$ -based primary productivity measurements at (a) P1 (SAZ-West) ( $r^2 = 0.98$ , slope = $0.79 \pm 0.06$ s.d., $n = 19$ , $p < 0.01$ ), (b) P2 (PFZ) ( $r^2 = 0.97$ , slope = $1.67 \pm 0.11$ s.d., $n = 6$ , $p < 0.01$ ), and (c) P3 (SAZ-East) ( $r^2 = 0.92$ , slope = $1.43 \pm 0.10$ s.d., $n = 14$ , $p < 0.01$ ). The dashed line represents the 1:1 relationship. P, Process station; SAZ, Subantarctic Zone; PFZ, Polar Frontal Zone. . . . .	24
3.1	SAZ-Sense voyage track and sampling locations superimposed on (a) MODIS-Aqua 9-km resolution chlorophyll $a$ concentration ( $\text{mg m}^{-3}$ ) and (b) Reynolds sea surface temperature ( $^{\circ}\text{C}$ ) daily composite image from 17 January to 20 February 2007. Red arrow indicates the approximate direction of the East Australian Current (EAC). Approximate locations of major fronts (STF, subtropical front; SAF-N, Sub-Antarctic front north; SAF-S, Sub-Antarctic front south; PF, polar front) are shown. Voyage track direction is shown by black arrows and black circles are the process stations (P1, process station 1; P2, process station 2; P3, process station 3). . . . .	32
3.2	Profiles of (a) mixed layer, $z_{\text{ML}}$ (m) and euphotic zone, $z_{\text{eu}}$ (m), mean surface PAR ( $\text{mol photons m}^{-2} \text{ d}^{-1}$ ) and mean PAR in the mixed layer, $E_{\text{ML}}$ ( $\text{mol photons m}^{-2} \text{ d}^{-1}$ ) at P1, P2, and P3, (b) log (base-10) transformed <i>in situ</i> PAR ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ), (c) chl $a$ ( $\mu\text{g L}^{-1}$ ) ( <a href="#">de Salas et al., 2011</a> ), (d) effective photochemical efficiency of PSII ( $F'_q/F'_m$ , dimensionless), and (d) functional absorption cross section of PSII ( $\sigma'_{\text{PSII}}$ , $\text{\AA}^2 \text{ quanta}^{-1}$ ). White lines in figure (b), (c), (d), and (e) are the mixed layer depths. . . . .	39
3.3	(a-c) Night-time values of maximum photochemical efficiency, $F_v/F_m$ (dimensionless) (black circle) and dissolved iron (red triangle) ( <a href="#">Lannuzel et al., 2011</a> ) at P1, P2 and a station near P3. (d-f) Functional absorption cross section of PSII, $\sigma_{\text{PSII}}$ ( $\text{\AA}^2 \text{ quanta}^{-1}$ ) (black square) and dissolved iron (red triangle) at P1, P2 and a station near P3. . . . .	40
3.4	Profiles of (a) light saturated photosynthesis rate, $P_{\text{max}}^{\text{B}}$ ( $\text{mg C (mg chl } a)^{-1} \text{ h}^{-1}$ ), (b) initial slope of the P-E curve, $\alpha^{\text{B}}$ ( $\text{mg C (mg chl } a)^{-1} \text{ h}^{-1} (\mu\text{mol photons m}^{-2} \text{ s}^{-1})^{-1}$ ), (c) saturation irradiance, $E_k$ ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) ( <a href="#">Westwood et al., 2011</a> ), (d) maximum rate of photosynthetic electron transport from charge separation to carbon fixation, $1/\tau_{\text{PSII}}$ ( $\text{ms}^{-1}$ ). . . . .	43
3.5	Profiles of (a) the sum of diadinoxanthin (dd) and diatoxanthin (dt) normalized to chl $a$ ( $\text{dd}+\text{dt}^*$ , dimensionless), (b) the de-epoxidation state of the xanthophyll cycle [ $\text{dt}/(\text{dd}+\text{dt})$ , dimensionless]. White lines indicate the mixed layer depth. . . . .	45
3.6	Chl $a$ -specific absorption coefficient of phytoplankton, $a_{\text{ph}}^*$ , at (a) 20 m, (b) 40m, (c) 60 m, (d) 80 m for P1, P2 and P3. . . . .	46
4.1	Surface photochemical efficiency of PSII ( $F_v/F_m$ , dimensionless) measurements along the Hobart-Dumont d'Urville transect in summer 2003. Colour lines are mean frontal positions from 2002 to 2009 based on <a href="#">Sokolov and Rintoul (2009a,b)</a> . STF, Subtropical Front; SAF-N, M, S, Subantarctic Front - North, Middle, South; PF-N,M,S, Polar Front - North, Middle, South; SACCF-N,S, Southern Antarctic Circumpolar Current Front - North, South; STZ, Subtropical Zone; SAZ, Subantarctic Zone; PFZ, Polar Frontal Zone; AZ, Antarctic Zone; SZ, Southern Zone. . . . .	57



- 4.2 Mean values of  $1^\circ$  latitude averaged sea surface (a) temperature (black) and salinity (blue), (b) nitrate (black) and chlorophyll (green), (c) phosphate, and (d) silicate from 2002 to 2009. All were *in situ* measurements except chlorophyll that were derived from SeaWiFS 9 km resolution 8-day averaged. Dashed lines indicate the standard deviation. Red lines are the front positions. 60
- 4.3 Distribution of surface (a) nitrate ( $\mu\text{M}$ ), (b) phosphate ( $\mu\text{M}$ ), and (c) silicate ( $\mu\text{M}$ ) along the Hobart-Dumont d'Urville transect from 2002 to 2009. Colour lines are mean frontal positions from north to south: STF (pink), SAZ (black), PF (blue), SACCF (red). . . . . 61
- 4.4 Distribution of surface  $F_v/F_m$  (dimensionless) along the Hobart-Dumont d'Urville transect from 2002 to 2009. a) Both day and nighttime measurements. b) Only nighttime measurements. Colour lines are mean frontal positions from north to south: STF (pink), SAZ (black), PF (blue), SACCF (red). . . . . 62
- 4.5 Averaged values of  $F_v/F_m$  (a) for the whole data sets, (b) in spring (Oct-Nov), (c) in summer (Dec-Jan), and (d) summer-early autumn (Feb-Mar). Black lines are day and night measurements and blue lines show only nighttime values. Dashed lines indicate the standard deviation. Red lines show the front positions. . . . . 63
- 4.6 Averaged diurnal pattern of  $F_v/F_m$  (black lines) and PAR (red lines). (a) All data, (b) STZ, (c) SAZ, (d) PFZ, (e) AZ, and (f) SZ. Dashed lines are the standard deviation. . . . . 64
- 4.7 Seasonal variations of diurnal  $F_v/F_m$  patterns in (a) STZ, (b) SAZ, (c) PFZ, (d) AZ, and (e) SZ. Dashed lines are standard deviation for each zone (see Fig. 4.5). (f) Distribution of  $F_v/F_m$  during the October-November 2003 transect with  $F_v/F_m$  in black and PAR in red. . . . . 65

## CHAPTER 1

### Introduction

#### 1 1.1 Marine primary producers

2 The emergence of phytoplankton more than 3 billion years ago altered the way our planet  
3 behaves and has had a profound effect on atmospheric chemistry and global climate. The  
4 relationship between photosynthetic organisms and global biogeochemical cycles is highly  
5 interlinked (*Falkowski and Raven, 2007*). The habitability of our planet depends on marine  
6 primary producers, where they contribute approximately 50% of global primary production,  
7 support nearly all life in the oceans and play a key role in the interlocking climate and  
8 biogeochemical systems.

9 If we are to fully understand the Earth system, it is important to be able to obtain  
10 accurate and precise estimates of the primary productivity of the oceans. The aggregate  
11 biomass of aquatic primary producers amounts to less than 1% of the total plant biomass  
12 on Earth (*Falkowski, 1994*). More than 95% of the carbon fixation in the oceans is the  
13 result of the photosynthetic activity of single-celled algae, the phytoplankton. Phytoplankton  
14 collectively fixes about 45 gigatons of organic carbon annually (*Falkowski et al., 1998*), which  
15 is equivalent in magnitude to global terrestrial primary production (*Behrenfeld et al., 2001*;  
16 *Cramer et al., 2001*). The turnover time of plants in marine ecosystems is about 0.02-0.06  
17 years compared to 9-20 years in terrestrial ecosystems (*Falkowski and Raven, 2007*). Changes  
18 in the oceanic primary production, linked to changes in the network of global biogeochemical  
19 cycles, can strongly influence atmospheric CO<sub>2</sub> on geological time scales. Understanding the  
20 changes in ocean primary productivity will help explain how oceanic biota responded to and  
21 are affected by natural climatic variability in the geological past, and how they will respond  
22 to anthropogenically influenced changes in coming decades (*Falkowski et al., 1998*).

## 1.2 The Southern Ocean

The Southern Ocean, which comprises approximately 10% of the world's oceans, is a major carbon sink for atmospheric CO<sub>2</sub> and plays an important role in controlling global climate. It also provides a unique environment that hosts a huge variety of marine organisms from single-celled phytoplankton to the largest mammals on Earth. The surface waters of the Southern Ocean are predicted to significantly change in response to climate change ([Sarmiento et al., 1998](#)). Climate-driven atmospheric and oceanic changes are changing the mixing rates of the ocean and thus modifying the nutrient supplies that are required for photosynthesis. These changes will affect phytoplankton species composition, dynamics and growth rates in many ways. Changes in oceanic phytoplankton communities will lead to changes in marine food webs and strongly influence the ability of Southern Ocean to absorb atmospheric CO<sub>2</sub>.

The Southern Ocean is characterised by dramatic gradients in physical and chemical properties, where, for example, sea surface temperatures can vary by up to 10 °C between its northern and southern boundaries ([Rintoul and Trull, 2001](#)). The hydrographic properties of the Southern Ocean are also characterized by extreme seasonal variability. This large variability makes the Southern Ocean a highly complex oceanographic region ([Rintoul and Trull, 2001](#)). Thus, marine organisms living in the Southern Ocean are exposed to these extreme environmental conditions where seawater temperature ranges from subzero to 20 °C. For phytoplankton, living in a such challenging environments requires fine tuning of their photosynthetic apparatus. The general seasonal pattern of phytoplankton blooms in the Southern Ocean (north of the ice edge) begins with winter, where cooling and strong winds extend the thermocline and enhance mixing of the upper ocean allowing nutrients to be mixed to the surface. In spring, increased sunlight, stratification and high nutrients concentration resulting from winter mixing in the euphotic zone lead to phytoplankton blooms. Apart from the physical challenges, phytoplankton in most of the Southern Ocean are also suffer from iron limitation. The s of macronutrients such as nitrate and phosphate are in excess all year round making the Southern Ocean the largest of the three major high nutrient

low chlorophyll (HNLC) regions in contemporary oceans (*de Baar et al., 2005*). Being the largest HNLC region, the Southern Ocean, has the greatest potential to affect atmospheric CO<sub>2</sub> concentration (*Sarmiento and Orr, 1991*).


### 1.3 Measuring primary production

Traditionally, primary production has been measured chemically. One method uses the evolution of O<sub>2</sub> in light bottles, which allows the estimation of net primary productivity and community respiration (*del Giorgio and Williams, 2005*). Another common technique uses radioactive <sup>14</sup>C. The advantage of the <sup>14</sup>C technique over the O<sub>2</sub> technique is the much greater sensitivity which permits measurement at much lower biomass concentration (*del Giorgio and Williams, 2005*). Since the introduction of the <sup>14</sup>C method in 1952 (*Steemann-Nielsen, 1952*), this technique has become the most common method to estimate primary productivity in oceanographic research, due to its comparative simplicity and sensitivity (*del Giorgio and Williams, 2005*).

Despite the accuracy and sensitivity offered by both techniques, a common disadvantage is that both techniques require the samples be incubated in a container, which will create artefacts. The artefacts are associated with isolating natural phytoplankton assemblages and containing them in bottles (*Eppley, 1980*), and differentiating between net and gross photosynthesis (*Bender et al., 1987*). Another disadvantage is that these techniques are time consuming and labourious and thus, limit the sampling rate. Furthermore, removal of phytoplankton from their natural environment limits the applicability of the results to the natural ecosystem. Assessment of primary production using satellite imagery has the potential to overcome some of these constraints and greatly improves spatial and temporal resolution (*Behrenfeld and Falkowski, 1997*). However, verification of these estimations with *in situ* measurements is always needed particularly at higher spatial and temporal resolution.

## 1.4 Fast repetition rate fluorometry

*In situ*, variable fluorescence-based measurements of phytoplankton photosynthesis potentially overcome many of the problems (e.g. artefacts from bottle incubation, labour intensive and time consuming) (Kolber and Falkowski, 1993). Variable fluorescence techniques provide valuable information of phytoplankton physiology in an instantaneous and *in situ* manner without the complication of artefacts from bottle incubations. One of the types of fluorometer that utilize this technique is the fast repetition rate (FRR) fluorometer (Fig. 1.1) (Kolber *et al.*, 1998).



1

Figure 1.1: Fast repetition rate fluorometer during deployment mode and measuring technique. The dual (light and dark) optical chambers contain series of blue light-emitting-diodes that provide subsaturating excitation flashes at 200 kHz repetition rate and fluorescence released from excited chlorophyll pigments will be detected by highly sensitive photomultiplier tube.

Using subsaturating (peak energy up to  $0.03 \text{ mol photons m}^{-2}\text{s}^{-1}$ ) excitation flashlets, the FRR fluorometer interrogate photosynthetic properties from light harvesting by photo-

85 system II (PSII) antennae to oxidation of the plastoquinone (PQ) and measures the initial  
 86 ( $F_o$ ), and maximal ( $F_m$ ) fluorescence yields resulting from the progressive closure of PSII  
 87 reaction centres and derives photosynthetic parameters such as the maximum photochemical  
 88 efficiency of PSII,  $F_v/F_m$ , and effective absorption cross-section of PSII,  $\sigma_{PSII}$  (Fig. 1.2).  
 89 FRR fluorescence yields were measured using a flash sequence consisting of a series of 100  
 90 subsaturation flashlets (1.1  $\mu s$  flash duration and 2.8  $\mu s$  inter flash period) (Fig. 1.3).




Figure 1.2: Simplified illustration of photosystem II. Light-harvesting pigments such as chlorophyll get excited after absorbing light energy (flashes). The absorbed light energy can then migrate to a reaction centre. If the reaction centre is open, minimum fluorescence  $F_o$  is released. If, at the instant of the photon absorption, the reaction centre is closed, the absorbed energy can be reradiated as fluorescence, increasing the fluorescence yield to  $F_m$ . Modified from (Kolber and Falkowski, 1993)

91 One of the major advantage of FRR fluorometry technique is the rapid measurement  
 92 rate of up to 1 second. This allows the photosynthetic parameters of the changing phyto-  
 93 plankton community to be monitored in near real time. The photosynthetic photosynthetic  
 94 parameters, such as the maximum photochemical efficiency of PSII ( $F_v/F_m$ ) and effective




Figure 1.3: Excitation flashes and relative fluorescence emission during a fast repetition rate fluorometry protocol. The excitation flashes consist of 100 subsaturation flashlets at  $1.1 \mu\text{s}$  flash duration and  $2.8 \mu\text{s}$  inter flash period.

absorption cross-section of PSII ( $\sigma_{\text{PSII}}$ ) are sensitive to environmental variables such as light and nutrients. For instance, a decline in  $F_v/F_m$  has been used as an indication of iron deficiency (*Behrenfeld et al., 1996*) in phytoplankton cells, a trace metal known to strongly limit primary productivity in up to 30% in the world's ocean (*de Baar et al., 2005*). In addition, when these photosynthetic parameters are combined with other environmental information, such as irradiance and biomass, estimation of primary production at fixed depths and continuous observations with a temporal resolution of seconds can be achieved (*Suggett et al., 2005*). Thus, FRR fluorometry offers unique *in situ* information on the photosynthetic competence and productivity of phytoplankton at a high temporal and spatial resolution, with minimal artefacts and allows rapid and real-time detection of change in phytoplankton.

## 1.5 Thesis objectives

The principle aim of this thesis is to investigate the coupling between the biological, chemical and physical processes that govern the physiological response and primary productivity of phytoplankton in the Southern Ocean, based on *in situ* observations measured with a FRR fluorometer and other oceanographic instruments. The main research themes of this thesis are:

- to investigate the relationship between FRR- and  $^{14}\text{C}$ -derived primary production in order to understand the reliability and applicability of FRR fluorometry to estimate primary productivity in the Southern Ocean.
- to assess how phytoplankton physiology in the Southern Ocean responds to the distinctive environmental variables.
- to investigate the seasonal and interannual variation in phytoplankton physiology in different water masses ranging from the subtropical to the polar waters.



1283

## Bibliography

- 1284 Alderkamp, A.-C., G. Kulk, A. G. J. Buma, R. J. W. Visser, G. L. Van Dijken, M. M. Mills, and  
 1285 K. R. Arrigo (2012), The effect of iron limitation on the photophysiology of *Phaeocystis antarctica*  
 1286 (Prymnesiophyceae) and *Fragilariopsis cylindrus* (Bacillariophyceae) under dynamic irradiance,  
 1287 *J. Phycol.*, *48*(1), 45–59, doi:10.1111/j.1529-8817.2011.01098.x.
- 1288 Allen, A. E., J. LaRoche, U. Maheswari, M. Lommer, N. Schauer, P. J. Lopez, G. Finazzi, A. R. Fernie,  
 1289 and C. Bowler (2008), Whole-cell response of the pennate diatom *Phaeodactylum tricornutum* to  
 1290 iron starvation, *Proceed. Natl. Acad. Sci.*, *105*(30), 10,438–10,443, doi:10.1073/pnas.0711370105.
- 1291 Arrigo, K., D. Worthen, A. Schnell, and M. Lizotte (1998), Primary production in Southern Ocean  
 1292 waters, *J. Geophys. Res.*, *103*(C8), 15,587–15,600, doi:10.1029/98JC00930.
- 1293 Arrigo, K., G. van Dijken, and S. Bushinsky (2008), Primary production in the Southern Ocean,  
 1294 1997–2006, *J. Geophys. Res.*, *113*, C08,004, doi:10.1029/2007JC004551.
- 1295 Bailey, S., et al. (2008), Alternative photosynthetic electron flow to oxygen in marine *Synechococcus*,  
 1296 *Biochim. Biophys. Acta Bioenerg.*, *1777*(3), 269–276, doi:10.1016/j.bbabbio.2008.01.002.
- 1297 Barlow, R., and R. Alberte (1985), Photosynthetic characteristics of phycoerythrin-containing marine  
 1298 *Synechococcus* spp. I. Responses to growth photon flux density, *Mar. Biol.*, *86*(1), 63–74, doi:  
 1299 10.1007/BF00392580.
- 1300 Behrenfeld, M. (2011), Uncertain future for ocean algae, *Nature Clim. Change*, *1*(1), 33–34, doi:  
 1301 10.1038/nclimate1069.
- 1302 Behrenfeld, M., and P. Falkowski (1997), A consumer’s guide to phytoplankton primary productivity  
 1303 models, *Limnol. Oceanogr.*, *42*(7), 1479–1491.
- 1304 Behrenfeld, M., et al. (2001), Biospheric primary production during an ENSO transition, *Science*,  
 1305 *291*(5513), 2594–2597, doi:10.1126/science.1055071.
- 1306 Behrenfeld, M. J., A. J. Bale, Z. S. Kolber, J. Aiken, and P. G. Falkowski (1996), Confirmation of  
 1307 iron limitation of phytoplankton photosynthesis in the equatorial Pacific Ocean, *Nature*, *383*(6600),  
 1308 508–511, doi:10.1038/383508a0.
- 1309 Behrenfeld, M. J., O. Prasil, Z. S. Kolber, M. Babin, and P. G. Falkowski (1998), Compensatory  
 1310 changes in Photosystem II electron turnover rates protect photosynthesis from photoinhibition,  
 1311 *Photosynth. Res.*, *58*, 259–268, doi:10.1023/A:1006138630573.
- 1312 Behrenfeld, M. J., K. Worthington, R. M. Sherrell, F. P. Chavez, P. Strutton, M. McPhaden, and  
 1313 D. M. Shea (2006), Controls on tropical Pacific Ocean productivity revealed through nutrient stress  
 1314 diagnostics, *Nature*, *442*(7106), 1025–1028, doi:10.1038/nature05083.
- 1315 Bender, M., et al. (1987), A comparison of four methods for determining planktonic community  
 1316 production, *Limnol. Oceanogr.*, pp. 1085–1098.
- 1317 Blain, S., et al. (2007), Effect of natural iron fertilization on carbon sequestration in the Southern  
 1318 Ocean, *Nature*, *446*(7139), 1070–1074, doi:10.1038/nature05700.
- 1319 Bowie, A. R., D. Lannuzel, T. A. Remenyi, T. Wagener, P. J. Lam, P. W. Boyd, C. Guieu, A. T.  
 1320 Townsend, and T. W. Trull (2009), Biogeochemical iron budgets of the Southern Ocean south of  
 1321 Australia: Decoupling of iron and nutrient cycles in the subantarctic zone by the summertime  
 1322 supply, *Global Biogeochem. Cycles*, *23*(4), GB4034, doi:10.1029/2009GB003500.
- 1323 Bowie, A. R., F. B. Griffiths, F. Dehairs, and T. W. Trull (2011a), Oceanography of the subantarctic  
 1324 and Polar Frontal Zones south of Australia during summer: Setting for the SAZ-Sense study,  
 1325 *Deep-Sea Res. II*, *58*(21–22), 2059 – 2070, doi:10.1016/j.dsr2.2011.05.033.

- 1326 Bowie, A. R., T. W. Trull, and F. Dehairs (2011b), Estimating the sensitivity of the subantarctic  
1327 zone to environmental change: The SAZ-Sense project, *Deep-Sea Res. II*, 58(21-22), 2051 – 2058,  
1328 doi:10.1016/j.dsr2.2011.05.034.
- 1329 Boyd, P., A. Crossley, G. DiTullio, F. Griffiths, D. Hutchins, B. Queguiner, P. Sedwick, and T. Trull  
1330 (2001), Control of phytoplankton growth by iron supply and irradiance in the subantarctic Southern  
1331 Ocean: Experimental results from the SAZ project, *J. Geophys. Res.*, 106(C12), 31,573–31,583.
- 1332 Boyd, P., R. Strzepek, F. Fu, and D. Hutchins (2010), Environmental control of open-ocean  
1333 phytoplankton groups: Now and in the future, *Limnol. Oceanogr.*, 55(3), 1353–1376, doi:  
1334 10.4319/lo.2010.55.3.1353.
- 1335 Boyd, P. W. (2002), Environmental factors controlling phytoplankton processes in the Southern  
1336 Ocean, *J. Phycol.*, 38(5), 844–861, doi:10.1046/j.1529-8817.2002.t01-1-01203.x.
- 1337 Boyd, P. W., and E. R. Abraham (2001), Iron-mediated changes in phytoplankton photo-  
1338 synthetic competence during SOIREE, *Deep-Sea Res. II*, 48(11-12), 2529–2550, doi:10.1016/  
1339 S0967-06450100007-8.
- 1340 Boyd, P. W., J. Aiken, and Z. Kolber (1997), Comparison of radiocarbon and fluorescence based  
1341 (pump and probe) measurements of phytoplankton photosynthetic characteristics in the Northeast  
1342 Atlantic Ocean, *Mar. Ecol. Prog. Ser.*, 149(1-3), 215–226, doi:10.3354/meps149215.
- 1343 Boyd, P. W., et al. (2000), A mesoscale phytoplankton bloom in the polar southern ocean stimulated  
1344 by iron fertilization, *Nature*, 407(6805), 695–702, doi:10.1038/35037500.
- 1345 Boyd, P. W., et al. (2007), Mesoscale iron enrichment experiments 1993-2005: Synthesis and future  
1346 directions, *Science*, 315(5812), 612–617, doi:10.1126/science.1131669.
- 1347 Brunet, C., and J. Lavaud (2010), Can the xanthophyll cycle help extract the essence of the microalgal  
1348 functional response to a variable light environment?, *J. Plankton Res.*, 32(12), 1609–1617, doi:  
1349 10.1093/plankt/fbq104.
- 1350 Butler, W. L. (1978), Energy distribution in the photochemical apparatus of photosynthesis, *Ann.*  
1351 *Rev. Plant Physiol.*, 29(1), 345–378, doi:10.1146/annurev.pp.29.060178.002021.
- 1352 Cassar, N., M. L. Bender, B. A. Barnett, S. Fan, W. J. Moxim, H. Levy, and B. Tilbrook (2007),  
1353 The Southern Ocean biological response to aeolian iron deposition, *Science*, 317(5841), 1067–1070,  
1354 doi:10.1126/science.1144602.
- 1355 Cheah, W., A. McMinn, F. B. Griffiths, K. J. Westwood, S. W. Wright, E. Molina, J. P. Webb, and  
1356 R. van den Enden (2011), Assessing Sub-Antarctic Zone primary productivity from fast repetition  
1357 rate fluorometry, *Deep-Sea Res. II*, 58(21-22), 2179 – 2188, doi:10.1016/j.dsr2.2011.05.023.
- 1358 Clementson, L., J. Parslow, A. Turnbull, D. McKenzie, and C. Rathbone (2001), Optical properties  
1359 of waters in the Australasian sector of the Southern Ocean, *J. Geophys. Res.*, 106(C12), 31,611–  
1360 31,625, doi:10.1029/2000JC000359.
- 1361 Coale, K. H., et al. (2004), Southern Ocean iron enrichment experiment: Carbon cycling in high- and  
1362 low-Si waters, *Science*, 304(5669), 408–414, doi:10.1126/science.1089778.
- 1363 Corno, G., R. M. Letelier, M. R. Abbott, and D. M. Karl (2006), Assessing primary production  
1364 variability in the North Pacific subtropical gyre: A comparison of fast repetition rate fluorometry  
1365 and  $^{14}\text{C}$  measurements, *J. Phycol.*, 42(1), 51–60, doi:10.1111/j.1529-8817.2006.00163.x.
- 1366 Cramer, W., et al. (2001), Global response of terrestrial ecosystem structure and function to  $\text{CO}_2$   
1367 and climate change: results from six dynamic global vegetation models, *Global Change Biol.*, 7(4),  
1368 357–373, doi:10.1046/j.1365-2486.2001.00383.x.

- Cubillos, J. C., S. W. Wright, G. Nash, M. F. de Salas, B. Griffiths, B. Tilbrook, A. Poisson, and G. M. Hallegraeff (2007), Calcification morphotypes of the coccolithophorid *Emiliania huxleyi* in the Southern Ocean: changes in 2001 to 2006 compared to historical data, *Mar. Ecol. Prog. Ser.*, *348*, 47–54, doi:10.3354/meps07058.
- de Baar, H. J. W., et al. (2005), Synthesis of iron fertilization experiments: From the iron age in the age of enlightenment, *J. Geophys. Res.*, *110*(C9), C09S16, doi:10.1029/2004JC002601.
- de Salas, M. F., R. Eriksen, A. T. Davidson, and S. W. Wright (2011), Protistan communities in the Australian sector of the Sub-Antarctic Zone during SAZ-Sense, *Deep-Sea Res. II*, *58*(21–22), 2135 – 2149, doi:10.1016/j.dsr2.2011.05.032.
- del Giorgio, P., and P. Williams (2005), *The global significance of respiration in aquatic ecosystems: from single cells to the biosphere*, 267 pp., Oxford University Press, USA.
- Demmig-Adams, B., and W. W. Adams (2006), Photoprotection in an ecological context: the remarkable complexity of thermal energy dissipation, *New Phytol.*, *172*(1), 11–21, doi:10.1111/j.1469-8137.2006.01835.x.
- El-Sayed, S. Z. (2005), History and evolution of primary productivity studies of the Southern Ocean, *Polar Biol.*, *28*(6), 423–438, doi:10.1007/s00300-004-0685-2.
- Eppley, R. (1972), Temperature and phytoplankton growth in the sea, *Fish. Bull.*, *70*(4), 1063–1085.
- Eppley, R. (1980), Estimating phytoplankton growth rates in the central oligotrophic oceans, in *Primary productivity in the sea*, vol. 31, edited by P. G. Falkowski, pp. 231–242, Plenum Press.
- Estevez-Blanco, P., P. Cermenio, M. Espineira, and E. Fernandez (2006), Phytoplankton photosynthetic efficiency and primary production rates estimated from fast repetition rate fluorometry at coastal embayments affected by upwelling (Rias Baixas, NW of Spain), *J. Plankton Res.*, *28*(12), 1153–1165, doi:10.1093/plankt/fbl046.
- Falkowski, P. (1994), The role of phytoplankton photosynthesis in global biogeochemical cycles, *Photosynth. Res.*, *39*(3), 235–258, doi:10.1007/BF00014586.
- Falkowski, P. G., and Z. Kolber (1995), Variations in chlorophyll fluorescence yields in phytoplankton in the world oceans, *Aust. Plant Physiol.*, *22*(2), 341–355, doi:10.1071/PP9950341.
- Falkowski, P. G., and J. A. Raven (2007), *Aquatic photosynthesis*, 2nd ed., Princeton University Press, Princeton, N. J.
- Falkowski, P. G., T. G. Owens, A. C. Ley, and D. C. Mauzerall (1981), Effects of growth irradiance levels on the ratio of reaction centers in 2 species of marine-phytoplankton, *Plant Physiol.*, *68*(4), 969–973, doi:10.1104/pp.68.4.969.
- Falkowski, P. G., Z. Dubinsky, and K. Wyman (1985), Growth-irradiance relationships in phytoplankton, *Limnol. Oceanogr.*, *30*(2), 311–321.
- Falkowski, P. G., Y. Fujita, A. Ley, and D. Mauzerall (1986), Evidence for cyclic electron flow around photosystem II in *Chlorellapyrenoidosa*, *Plant Physiol.*, *81*(1), 310–312, doi:10.1104/pp.81.1.310.
- Falkowski, P. G., R. T. Barber, and V. Smetacek (1998), Biogeochemical controls and feedbacks on ocean primary production, *Science*, *281*(5374), 200–206, doi:10.1126/science.281.5374.200.
- Flameling, I. A., and J. Kromkamp (1997), Photoacclimation of *Scenedesmus protuberans* (Chlorophyceae) to fluctuating irradiances simulating vertical mixing, *J. Plankton Res.*, *19*(8), 1011–1024, doi:10.1093/plankt/19.8.1011.
- Fujiki, T., T. Suzue, H. Kimoto, and T. Saino (2007), Photosynthetic electron transport in *Dunaliella tertiolecta* (Chlorophyceae) measured by fast repetition rate fluorometry: relation to carbon assimilation, *J. Plankton Res.*, *29*(2), 199–208, doi:10.1093/plankt/fbm007.

- Gallegos, C. L., and T. Platt (1985), Vertical advection of phytoplankton and productivity estimates: A dimensional analysis, *Mar. Ecol. Prog. Ser.*, 26(1-2), 125–134.
- Gervais, F., U. Riebesell, and M. Y. Gorbunov (2002), Changes in primary productivity and chlorophyll a in response to iron fertilization in the Southern Polar Frontal Zone, *Limnol. Oceanogr.*, 47(5), 1324–1335.
- Goss, R., and T. Jakob (2010), Regulation and function of xanthophyll cycle-dependent photoprotection in algae, *Photosynth. Res.*, 106, 103–122, doi:10.1007/s11120-010-9536-x.
- Griffiths, F. B., T. S. Bates, P. K. Quinn, L. A. Clementson, and J. S. Parslow (1999), Oceanographic context of the First Aerosol Characterization Experiment (ACE 1): A physical, chemical, and biological overview, *J. Geophys. Res.*, 104(D17), 21,649–21,671, doi:10.1029/1999JD900386.
- Gruber, N., et al. (2009), Oceanic sources, sinks, and transport of atmospheric CO<sub>2</sub>, *Global Biogeochem. Cycles*, 23(1), GB1005, doi:10.1029/2008GB003349.
- Heber, U. (2002), Irrungen, Wirrungen? The Mehler reaction in relation to cyclic electron transport in C3 plants, *Photosynth. Res.*, 73(1-3), 223–231, doi:10.1023/A:1020459416987.
- Herraiz-Borreguero, L., and S. R. Rintoul (2011), Regional circulation and its impact on upper ocean variability south of Tasmania, *Deep-Sea Res. II*, 58(21-22), 2071 – 2081, doi:10.1016/j.dsr2.2011.05.022.
- Higgins, H. W., S. W. Wright, and L. Schlüter (2011), Quantitative interpretation of chemotaxonomic pigment data, in *Phytoplankton pigments: Characterization, chemotaxonomy and applications in oceanography*, edited by S. Roy, E. S. Egeland, G. Johnsen, and C. A. Llewellyn, chap. 6, pp. 257–313, Cambridge University Press.
- Hill, K. L., S. R. Rintoul, R. Coleman, and K. R. Ridgway (2008), Wind forced low frequency variability of the East Australia Current, *Geophys. Res. Lett.*, 35(8), L08,602, doi:10.1029/2007GL032912.
- Holeton, C. L., F. Nedelec, R. Sanders, L. Brown, C. M. Moore, D. P. Stevens, K. J. Heywood, P. J. Statham, and C. H. Lucas (2005), Physiological state of phytoplankton communities in the Southwest Atlantic sector of the Southern Ocean, as measured by fast repetition rate fluorometry, *Polar Biol.*, 29(1), 44–52, doi:10.1007/s00300-005-0028-y.
- Hutchins, D. A., P. N. Sedwick, G. R. DiTullio, P. W. Boyd, B. Queguiner, F. B. Griffiths, and C. Crossley (2001), Control of phytoplankton growth by iron and silicic acid availability in the subantarctic Southern Ocean: Experimental results from the SAZ Project, *J. Geophys. Res.*, 106(C12), 31,559–31,572, doi:10.1029/2000JC000333.
- Jakob, T., R. Goss, and C. Wilhelm (2001), Unusual ph-dependence of diadinoxanthin de-epoxidase activation causes chlororespiratory induced accumulation of diatoxanthin in the diatom *Phaeodactylum tricornutum*, *J. Plant Physiol.*, 158(3), 383 – 390, doi:10.1078/0176-1617-00288.
- Kana, T. M. (1992), Relationship between photosynthetic oxygen cycling and carbon assimilation in *Synechococcus* WH7803 (Cyanophyta), *J. Phycol.*, 28(3), 304–308, doi:10.1111/j.0022-3646.1992.00304.x.
- Kana, T. M. (1993), Rapid oxygen cycling in *Trichodesmium thiebautii*, *Limnol. Oceanogr.*, 38(1), 18–24.
- Koblížek, M., D. Kaftan, and L. Nedbal (2001), On the relationship between the non-photochemical quenching of the chlorophyll fluorescence and the Photosystem II light harvesting efficiency. A repetitive flash fluorescence induction study, *Photosynth. Res.*, 68(2), 141–152, doi:10.1023/A:1011830015167.
- Kolber, Z., and P. G. Falkowski (1993), Use of active fluorescence to estimate phytoplankton photosynthesis in situ, *Limnol. Oceanogr.*, 38(8), 1646–1665.

- 1458 Kolber, Z. S., R. T. Barber, K. H. Coale, S. E. Fitzwater, R. M. Greene, K. S. Johnson, S. Lindley, and  
1459 P. G. Falkowski (1994), Iron limitation of phytoplankton photosynthesis in the equatorial Pacific  
1460 Ocean, *Nature*, 371(6493), 145–149, doi:10.1038/371145a0.
- 1461 Kolber, Z. S., O. Prasil, and P. G. Falkowski (1998), Measurements of variable chlorophyll fluorescence  
1462 using fast repetition rate techniques: defining methodology and experimental protocols, *Biochim.*  
1463 *Biophys. Acta*, 1367(1-3), 88–106, doi:10.1016/S0005-2728(98)00135-2.
- 1464 Korb, R. E., M. J. Whitehouse, A. Atkinson, and S. E. Thorpe (2008), Magnitude and maintenance  
1465 of the phytoplankton bloom at South Georgia: a naturally iron-replete environment, *Mar. Ecol.*  
1466 *Prog. Ser.*, 368, 75–91, doi:10.3354/meps07525.
- 1467 Kosakowska, A., J. Lewandowska, J. Stoń, and K. Burkiewicz (2004), Qualitative and quantita-  
1468 tive composition of pigments in *Phaeodactylum tricornutum* (Bacillariophyceae) stressed by iron,  
1469 *Biometals*, 17(1), 45–52, doi:10.1023/A:1024452802005.
- 1470 Lannuzel, D., V. Schoemann, J. de Jong, J.-L. Tison, and L. Chou (2007), Distribution and bio-  
1471 geochemical behaviour of iron in the East Antarctic sea ice, *Mar. Chem.*, 106(1-2), 18–32, doi:  
1472 10.1016/j.marchem.2006.06.010.
- 1473 Lannuzel, D., A. R. Bowie, T. Remenyi, P. Lam, A. Townsend, E. Ibanmi, E. Butler, T. Wagener,  
1474 and V. Schoemann (2011), Distributions of dissolved and particulate iron in the sub-Antarctic  
1475 and Polar Frontal Southern Ocean (Australian sector), *Deep-Sea Res. II*, 58(21-22), 2094–2112,  
1476 doi:10.1016/j.dsr2.2011.05.027.
- 1477 Lavaud, J., and P. G. Kroth (2006), In diatoms, the transthylakoid proton gradient regulates the  
1478 photoprotective non-photochemical fluorescence quenching beyond its control on the xanthophyll  
1479 cycle, *Plant Cell Physiol.*, 47(7), 1010–1016, doi:10.1093/pcp/pcj058.
- 1480 Laws, E. A. (1991), Photosynthetic quotients, new production and net community production in the  
1481 open ocean, *Deep-Sea Res. I*, 38(1), 143–167, doi:10.1016/0198-01499190059-O.
- 1482 Lewis, M. R., and J. C. Smith (1983), A small volume, short-Incubation-time method for measurement  
1483 of photosynthesis as a function of incident irradiance, *Mar. Ecol. Prog. Ser.*, 13(1), 99–102.
- 1484 Li, W. K. W. (1994), Primary production of prochlorophytes, cyanobacteria, and eukaryotic ultraphy-  
1485 toplanktons - measurements from flow cytometric sorting, *Limnol. Oceanogr.*, 39(1), 169–175.
- 1486 Li, Z., S. Wakao, B. B. Fischer, and K. K. Niyogi (2009), Sensing and responding to excess light,  
1487 *Annu. Rev. Plant Biol.*, 60(1), 239–260, doi:10.1146/annurev.arplant.58.032806.103844.
- 1488 Lovenduski, N. S., and N. Gruber (2005), Impact of the Southern Annular Mode on Southern Ocean  
1489 circulation and biology, *Geophys. Res. Lett.*, 32(11), L11,603, doi:10.1029/2005gl022727.
- 1490 Macedo, M. F., P. Duarte, and J. G. Ferreira (2002), The influence of incubation periods on  
1491 photosynthesis-irradiance curves, *J. Exp. Mar. Biol. Ecol.*, 274(2), 101–120, 10.1016/S0022-  
1492 0981(02)00202-2.
- 1493 Marra, J. (2002), Approaches to the measurement of plankton production, in *Phytoplankton produc-*  
1494 *tivity*, edited by P. Williams, D. Thomas, and C. Reynolds, pp. 78–108, Blackwell Science Ltd.,  
1495 Oxford, doi:10.1002/9780470995204.ch4.
- 1496 McLeod, D. J., G. M. Hallegraeff, G. W. Hosie, and A. J. Richardson (2012), Climate-driven range  
1497 expansion of the red-tide dinoflagellate *noctiluca scintillans* into the southern ocean, *Journal of*  
1498 *Plankton Research*, 34(4), 332–337, doi:10.1093/plankt/fbr112.
- 1499 McNeil, B. I., B. Tilbrook, and R. J. Matear (2001), Accumulation and uptake of anthropogenic CO<sub>2</sub>  
1500 in the Southern Ocean, south of Australia between 1968 and 1996, *J. Geophys. Res.*, 106(C12),  
1501 31,431–31,445, doi:10.1029/2000jc000331.

- Melrose, D. C., C. A. Oviatt, J. E. O'Reilly, and M. S. Berman (2006), Comparisons of fast repetition rate fluorescence estimated primary production and  $^{14}\text{C}$  uptake by phytoplankton, *Mar. Ecol. Prog. Ser.*, *311*, 37–46, doi:10.3354/meps311037.
- Metzl, N., B. Tilbrook, and A. Poisson (1999), The annual  $f\text{CO}_2$  cycle and the air-sea  $\text{CO}_2$  flux in the sub-Antarctic Ocean, *Tellus Ser. B.*, *51*(4), 849–861, doi:10.1034/j.1600-0889.1999.t01-3-00008.x.
- Metzl, N., C. Brunet, A. Jabaud-Jan, A. Poisson, and B. Schauer (2006), Summer and winter air-sea  $\text{CO}_2$  fluxes in the Southern Ocean, *Deep-Sea Res. I*, *53*(9), 1548–1563, doi:10.1016/j.dsr.2006.07.006.
- Mongin, M., R. Matear, and M. Chamberlain (2011), Seasonal and spatial variability of remotely sensed chlorophyll and physical fields in the saz-sense region, *Deep Sea Research Part II: Topical Studies in Oceanography*, *58*(21-22), 2082 – 2093, doi:10.1016/j.dsr2.2011.06.002.
- Moore, C. M., et al. (2003), Physical controls on phytoplankton physiology and production at a shelf sea front: a fast repetition-rate fluorometer based field study, *Mar. Ecol. Prog. Ser.*, *259*, 29–45, doi:10.3354/meps259029.
- Moore, C. M., D. J. Suggett, A. E. Hickman, Y.-N. Kim, J. F. Tweddle, J. Sharples, R. J. Geider, and P. M. Holligan (2006), Phytoplankton photoacclimation and photoadaptation in response to environmental gradients in a shelf sea, *Limnol. Oceanogr.*, *51*(2), 936–949.
- Moore, J. K., and M. R. Abbott (2000), Phytoplankton chlorophyll distributions and primary production in the Southern Ocean, *J. Geophys. Res.*, *105*(C12), 28,709–28,722, doi:10.1029/1999JC000043.
- Moseley, J. L., T. Allinger, S. Herzog, P. Hoerth, E. Wehinger, S. Merchant, and M. Hippler (2002), Adaptation to Fe-deficiency requires remodeling of the photosynthetic apparatus, *EMBO J*, *21*(24), 6709–6720, doi:10.1093/emboj/cdf666.
- Ogren, W. L. (1984), Photorespiration: Pathways, regulation, and modification, *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, *35*, 415–442.
- Olaizola, M., and H. Y. Yamamoto (1994), Short-term response of the diadinoxanthin cycle and fluorescence yield to high irradiance in *chaetoceros muelleri* (bacillariophyceae), *J. Phycol.*, *30*(4), 606–612, doi:10.1111/j.0022-3646.1994.00606.x.
- Olson, R. J., H. M. Sosik, A. M. Chekalyuk, and A. Shalapyonok (2000), Effects of iron enrichment on phytoplankton in the Southern Ocean during late summer: active fluorescence and flow cytometric analyses, *Deep-Sea Res. II*, *47*(15-16), 3181–3200, doi:10.1016/S0967-06450000064-3.
- Partensky, F., W. R. Hess, and D. Vaulot (1999), *Prochlorococcus*, a marine photosynthetic prokaryote of global significance, *Microbiol. Mol. Biol. Rev.*, *63*(1), 106–127.
- Platt, T., C. L. Gallegos, and W. G. Harrison (1980), Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton, *J. Mar. Res.*, *38*(4), 687–701.
- Pollard, R. T., et al. (2009), Southern Ocean deep-water carbon export enhanced by natural iron fertilization, *Nature*, *457*(7229), 577–581, doi:10.1038/nature07716.
- Prasil, O., Z. Kolber, J. A. Berry, and P. G. Falkowski (1996), Cyclic electron flow around photosystem II in vivo, *Photosynth. Res.*, *48*(3), 395–410, doi:10.1007/BF00029472.
- Raateoja, M., J. Seppala, and H. Kuosa (2004), Bio-optical modelling of primary production in the SW Finnish coastal zone, Baltic Sea: fast repetition rate fluorometry in Case 2 waters, *Mar. Ecol. Prog. Ser.*, *267*, 9–26, doi:10.3354/meps267009.
- Raateoja, M., B. G. Mitchell, H. Wang, and E. Olivo (2009), Effect of water column light gradient on phytoplankton fluorescence transients, *Mar. Ecol. Prog. Ser.*, *376*, 85–101, doi:10.3354/meps07759.

- 1546 Raven, J. A., and R. J. Geider (1988), Temperature and algal growth, *New Phytol.*, *110*(4), 441–461,  
1547 doi:10.1111/j.1469-8137.1988.tb00282.x.
- 1548 Reay, D. S., J. Priddle, D. B. Nedwell, M. J. Whitehouse, J. C. Ellis-Evans, C. Deubert, and D. P.  
1549 Connelly (2001), Regulation by low temperature of phytoplankton growth and nutrient uptake in  
1550 the Southern Ocean, *Mar. Ecol. Prog. Ser.*, *219*, 51–64, doi:10.3354/meps219051.
- 1551 Ridgway, K. R. (2007), Long-term trend and decadal variability of the southward penetration of the  
1552 East Australian Current, *Geophys. Res. Lett.*, *34*(13), L13,613, doi:10.1029/2007gl030393.
- 1553 Rintoul, S. R., and T. W. Trull (2001), Seasonal evolution of the mixed layer in the Subantarctic  
1554 Zone south of Australia, *J. Geophys. Res.*, *106*(C12), 31,447–31,462, doi:10.1029/2000jc000329.
- 1555 Rose, J. M., et al. (2009), Synergistic effects of iron and temperature on Antarctic phytoplankton and  
1556 microzooplankton assemblages, *Biogeosciences*, *6*(12), 3131–3147, doi:10.5194/bg-6-3131-2009.
- 1557 Sarmiento, J., and J. Orr (1991), Three-Dimensional Simulations of the Impact of Southern Ocean  
1558 Nutrient Depletion on Atmospheric CO<sub>2</sub> and Ocean Chemistry, *Limnol. Oceanogr.*, *36*(8), 1928–  
1559 1950.
- 1560 Sarmiento, J. L., T. M. C. Hughes, R. J. Stouffer, and S. Manabe (1998), Simulated response of the  
1561 ocean carbon cycle to anthropogenic climate warming, *Nature*, *393*(6682), 245–249, doi:10.1038/  
1562 30455.
- 1563 Sarmiento, J. L., N. Gruber, M. A. Brzezinski, and J. P. Dunne (2004), High-latitude controls of  
1564 thermocline nutrients and low latitude biological productivity, *Nature*, *427*(6969), 56–60, doi:  
1565 10.1038/nature02127.
- 1566 Sarmiento, J. L., M. Gloor, N. Gruber, C. Beaulieu, A. R. Jacobson, S. E. Mikaloff Fletcher, S. Pacala,  
1567 and K. Rodgers (2010), Trends and regional distributions of land and ocean carbon sinks, *Biogeo-*  
1568 *sciences*, *7*(8), 2351–2367, doi:10.5194/bg-7-2351-2010.
- 1569 Sedwick, P., A. Bowie, and T. Trull (2008), Dissolved iron in the Australian sector of the Southern  
1570 Ocean (CLIVAR SR3 section): Meridional and seasonal trends, *Deep-Sea Res. I*, *55*(8), 911–925,  
1571 doi:10.1016/j.dsr.2008.03.011.
- 1572 Sedwick, P. N., P. R. Edwards, D. J. Mackey, F. B. Griffiths, and J. S. Parslow (1997), Iron and  
1573 manganese in surface waters of the Australian subantarctic region, *Deep-Sea Res. I*, *44*(7), 1239–  
1574 1253, doi:10.1016/S0967-06379700021-6.
- 1575 Smyth, T. J., K. L. Pemberton, J. Aiken, and R. J. Geider (2004), A methodology to determine  
1576 primary production and phytoplankton photosynthetic parameters from Fast Repetition Rate Flu-  
1577 orometry, *J. Plankton Res.*, *26*(11), 1337–1350, doi:10.1093/plankt/fbh124.
- 1578 Sokolov, S., and S. R. Rintoul (2002), Structure of Southern Ocean fronts at 140°E, *J. Mar. Syst.*,  
1579 *37*(1-3), 151–184, doi:10.1016/S0924-79630200200-2.
- 1580 Sokolov, S., and S. R. Rintoul (2003), Subsurface structure of interannual temperature anomalies  
1581 in the Australian sector of the Southern Ocean, *J. Geophys. Res.*, *108*(C9), 3285, doi:10.1029/  
1582 2002jc001494.
- 1583 Sokolov, S., and S. R. Rintoul (2007), On the relationship between fronts of the Antarctic Circumpolar  
1584 Current and surface chlorophyll concentrations in the Southern Ocean, *J. Geophys. Res.*, *112*(C7),  
1585 C07,030, doi:10.1029/2006jc004072.
- 1586 Sokolov, S., and S. R. Rintoul (2009a), Circumpolar structure and distribution of the Antarctic  
1587 Circumpolar Current fronts: 1. Mean circumpolar paths, *J. Geophys. Res.*, *114*(C11), C11,018,  
1588 doi:10.1029/2008jc005108.
- 1589 Sokolov, S., and S. R. Rintoul (2009b), Circumpolar structure and distribution of the Antarctic  
1590 Circumpolar Current fronts: 2. Variability and relationship to sea surface height, *J. Geophys. Res.*,  
1591 *114*(C11), C11,019, doi:10.1029/2008jc005248.

- Sosik, H. M., and R. J. Olson (2002), Phytoplankton and iron limitation of photosynthetic efficiency in the Southern Ocean during late summer, *Deep-Sea Res. I*, *49*(7), 1195 – 1216, doi:10.1016/S0967-0637(02)00015-8.
- Steemann-Nielsen, E. (1952), The use of radioactive carbon ( $^{14}\text{C}$ ) for measuring organic production in the sea, *J. Cons. Int. Explor. Mer*, *18*, 117–140.
- Strutton, P. G., F. B. Griffiths, R. L. Waters, S. W. Wright, and N. L. Bindoff (2000), Primary productivity off the coast of East Antarctica (80-150°E): January to March 1996, *Deep-Sea Res. II*, *47*(12-13), 2327–2362, doi:10.1016/S0967-06450000028-X.
- Strzepek, R. F., and P. J. Harrison (2004), Photosynthetic architecture differs in coastal and oceanic diatoms, *Nature*, *431*(7009), 689–92, doi:10.1038/nature02954.
- Suggett, D., G. Kraay, P. Holligan, M. Davey, J. Aiken, and R. Geider (2001), Assessment of photosynthesis in a spring cyanobacterial bloom by use of a fast repetition rate fluorometer, *Limnol. Oceanogr.*, *46*(4), 802–810.
- Suggett, D. J., C. M. Moore, K. Oxborough, and R. J. Geider (2005), *Fast repetition rate (FRR) chlorophyll a fluorescence induction measurements*, Chelsea Technologies Group, Surrey.
- Suggett, D. J., C. M. Moore, E. Marañón, C. Omachi, R. A. Varela, J. Aiken, and P. M. Holligan (2006), Photosynthetic electron turnover in the tropical and subtropical Atlantic Ocean, *Deep-Sea Res. II*, *53*(14-16), 1573–1592, doi:10.1016/j.dsr2.2006.05.014.
- Suggett, D. J., C. M. Moore, A. E. Hickman, and R. G. Geider (2009), Interpretation of fast repetition rate (FRR) fluorescence: signatures of phytoplankton community structure versus physiological state, *Mar. Ecol. Prog. Ser.*, *376*, 1 – 9, doi:10.3354/meps07830.
- Sullivan, C. W., K. R. Arrigo, C. R. McClain, J. C. Comiso, and J. Firestone (1993), Distributions of phytoplankton blooms in the Southern Ocean, *Science*, *262*(5141), 1832–1837, doi:10.1126/science.262.5141.1832.
- Sunda, W. G., and S. A. Huntsman (1997), Interrelated influence of iron, light and cell size on marine phytoplankton growth, *Nature*, *390*(6658), 389–392, doi:10.1038/37093.
- Suzuki, K., H. Liu, T. Saino, H. Obata, M. Takano, K. Okamura, Y. Sohrin, and Y. Fujishima (2002), East-west gradients in the photosynthetic potential of phytoplankton and iron concentration in the subarctic pacific ocean during early summer, *Limnology and oceanography*, pp. 1581–1594, doi:10.4319/lo.2002.47.6.1581.
- Takahashi, T., et al. (2009), Climatological mean and decadal change in surface ocean  $p\text{CO}_2$ , and net sea-air  $\text{CO}_2$  flux over the global oceans, *Deep-Sea Res. II*, *56*(8-10), 554–577.
- Thompson, D. W. J., and J. M. Wallace (2000), Annular modes in the extratropical circulation. Part I: Month-to-month variability, *J. Clim.*, *13*(5), 1000–1016, doi:10.1175/1520-0442(2000)013<1000:AMITEC>2.0.CO;2.
- Tilzer, M. M., m. Elbrächter, W. W. Gieskes, and B. Beese (1986), Light-temperature interactions in the control of photosynthesis in Antarctic phytoplankton, *Polar Biol.*, *5*, 105–111, doi:10.1007/BF00443382.
- Trull, T., S. R. Rintoul, M. Hadfield, and E. R. Abraham (2001), Circulation and seasonal evolution of polar waters south of Australia: implications for iron fertilization of the Southern Ocean, *Deep-Sea Res. II*, *48*(11-12), 2439–2466, doi:10.1016/S0967-0645(01)00003-0.
- Vaillancourt, R. D., R. N. Sambrotto, S. Green, and A. Matsuda (2003), Phytoplankton biomass and photosynthetic competency in the summertime Mertz Glacier Region of East Antarctica, *Deep-Sea Res. II*, *50*(8-9), 1415–1440, doi:10.1016/S0967-06450300077-8.



- 1636 van de Poll, W., P. Janknegt, M. van Leeuwe, R. Visser, and A. Buma (2009), Excessive irradiance  
1637 and antioxidant responses of an antarctic marine diatom exposed to iron limitation and to dynamic  
1638 irradiance, *J. Photoch. Photobio. B*, *94*(1), 32–37, doi:10.1016/j.jphotobiol.2008.09.003.
- 1639 van de Poll, W. H., and A. G. J. Buma (2009), Does ultraviolet radiation affect the xanthophyll cycle  
1640 in marine phytoplankton?, *Photoch. Photobio. Sci.*, *8*, 1295–1301, doi:10.1039/B904501E.
- 1641 van de Poll, W. H., R. J. W. Visser, and A. G. J. Buma (2007), Acclimation to a dynamic irra-  
1642 diance regime changes excessive irradiance sensitivity of *Emiliana huxleyi* and *Thalassiosira*  
1643 *weissflogii*, *Limnol. Oceanogr.*, *52*(4), 1430–1438, doi:10.4319/lo.2007.52.4.1430.
- 1644 Venables, H., and C. M. Moore (2010), Phytoplankton and light limitation in the Southern  
1645 Ocean: Learning from high-nutrient, high-chlorophyll areas, *J. Geophys. Res.*, *115*, doi:10.1029/  
1646 2009JC005361.
- 1647 Westwood, K. J., F. B. Griffiths, J. P. Webb, and S. W. Wright (2011), Primary production in the  
1648 Sub-Antarctic and Polar Frontal Zones south of Tasmania, Australia; SAZ-Sense survey, 2007,  
1649 *Deep-Sea Res. II*, *58*(21-22), 2162 – 2178, doi:10.1016/j.dsr2.2011.05.017.
- 1650 Williams, P. J. L., and D. Lefevre (1996), Algal  $^{14}\text{C}$  and total carbon metabolisms. 2. Experimental  
1651 observations with the diatom *Skeletonema costatum*, *J. Plankton Res.*, *18*(10), 1941–1959, doi:  
1652 10.1093/plankt/18.10.1961.
- 1653 Wolter, K., and M. Timlin (1998), Measuring the strength of ENSO events: How does 1997/98 rank?,  
1654 *Weather*, *53*, 315–323.
- 1655 Wright, S. W., D. P. Thomas, H. J. Marchant, H. W. Higgins, M. D. Mackey, and D. J. Mackey  
1656 (1996), Analysis of phytoplankton of the Australian sector of the Southern Ocean: Compar-  
1657 isons of microscopy and size frequency data with interpretations of pigment HPLC data using  
1658 the 'CHEMTAX' matrix factorisation program, *Mar. Ecol. Prog. Ser.*, *144*(1-3), 285–298, doi:  
1659 10.3354/meps144285.
- 1660 Wright, S. W., R. L. van den Enden, I. Pearce, A. T. Davidson, F. J. Scott, and K. J. Westwood (2010),  
1661 Phytoplankton community structure and stocks in the Southern Ocean (30-80°E) determined by  
1662 CHEMTAX analysis of HPLC pigment signatures, *Deep-Sea Res. II*, *57*(9-10), 758–778, doi:10.  
1663 1016/j.dsr2.2009.06.015.
- 1664 Zapata, M., F. Rodríguez, and J. Garrido (2000), Separation of chlorophylls and carotenoids from  
1665 marine phytoplankton: a new HPLC method using a reversed phase C8 column and pyridine-  
1666 containing mobile phases, *Mar. Ecol. Prog. Ser.*, *195*, 29–45, doi:10.3354/meps195029.